

55. Türkiye Jeoloji Kurultayı
55th Geological Congress of Turkey

KIRŞEHİR KUZEYİNDEKİ MAGMATİK KAYAÇLAR VE MADEN YATAKLARI

Ramazan DOĞAN, Emrah AYAZ, LDeniz ÖNENÇ, Kamil UÇAR, Okan ZİMİTOĞLU
MTA Genel Müdürlüğü, 06520 Ankara

Kırşehir kuzeyindeki magmatik kayaçlar ve bunlarla ilgili maden yatakları 2001 yılında MTA'nın Âsıdık Magmatizmaya Bağlı Maden Yatakları Aramaları Projesi çerçevesinde çalışılmış ve Yozgat güneyinde ortaya konulan magmatik gruplara ek olarak beş ayrı magmatik topluluğun varlığı saptanmıştır. Bölgedeki bütün magmatik kayaçlar yaşlıdan gence doğru şu şekilde sıralanabilir: 1) Meta-bazik kayaçlar, 2) Lökogranitler, 3) Cefalik dağı granitleri, 4) Gabro-bazaltlar, 5) Gabro-monzonitler, 6) Granodiyorit-granitler, 7) Siyenitler, 8) Çok alkali kayaçlar, 9) Tersiyer yaşlı volkanik kayaçlar,

Meta-bazik kayaçlar yeşil şist metamorfizmasına uğramış gabro, bazalt, volkano-sedimanter kayaçlardan ve az miktarda asıdık volkanik kayaçlardan oluşmuştur* Balıh-Sulakyurt arasında yaygın olarak mostra veren meta-bazik kayaçlar içinde bazı küçük demir madenlerine rastlanmıştır, Lökogranitler genellikle KD-GB yönünde uzanan zonlar boyunca meta-bazik kayaçlar içine sokulmuşlardır. Aplit ve pegmatit stok ve daykları Yozgat'a göre daha az gelişmiştir. Lökogranitler ayrıca Keskinin batısında ve Behrekdağı'nın KD-sunda da yüzeylenir,

Kaman'n güneyindeki Cefalik dağı granitleri güneyde migmatitik karakter göstermektedir. Plutonun bilhassa kenar kısımlarında kayaç içindeki mineral dağılımı yer yer çok heterojen olup; kayaç, etrafındaki yüksek dereceli metamorfik kayaçlara benzer yapısal unsurlar içerir, Savcılıbeyit yakınlarındaki altın zuhurlarının da metamorfizme sırasında oluşan segregasyon kuvarları ile birlikte meydana geldikleri sanılmaktadır,

Gabro-bazalt grubu magmatik kayaçlar Kaman'ın KB'sından geçen KD-GB yönlü ve Akçakentten geçen KDD-GBB yönlü graben havzaları içinde yer alır. Keskinen geçen KB-GD yönlü havza iyi gelişmemiştir. Grabenin kenar zonlarında gabro ve monzonit daykları, iç kısımlarında ise bazalt ve radyoarit içeren volkano-sedimanter kayaçlar görülür. Kaşmağa yakınlarında işletilmiş birkaç Mn ocağı ve yüksek Zn anomalisi bu ortam için karakteristik cevherleşmelerdir. Graben yapısının Cefalik Dağı granitlerini kesmesi ve bu veya bir sonraki grup üyesi gabroların lökogranitleri anklav olarak içermesi, gruplar arasındaki bağıl yaş ilişkilerinin saptanmasında önemli olan unsurlardandır,

Gabro-monzonit grubu kayaçlar Baranadağ'da, Akçakentte ve Dinek dağı'nın güneyinde görülürler. Sorgun'un güneyinde Kerkenez dağında olduğu gibi, dıştan içe doğru gabro-diyorit-monzodiyorit-porfiritik monzonit-porfiritik kuvar monzonit normal zonlaşması buralarda da gelişmiştir, Bayındır ve Akpınar güneyinde ve Akçakent Madentepe'de olduğu gibi gabrolar içinde bazen dasit porfiri daykları ve silisleşme ile birlikte Fe cevherleşmeleri oluşmuştur. Yüzey ve/veya hidrotermal

55, Türkiye jeoloji Kurultayı
55th Geological Congress of Turkey

alterasyona uğramış porfiritik monzonitlerde K-feldispat fenokristalleri Kievi'nde olduğu gibi, bazen basit bir tesiste kolaylıkla zenginleştirilebilmektedir.

Granodiyorit-granit grubu kayalar Yozgat güneyindeklere göre kısa aralıklarda daha farklı kompozisyonlar sergilemektedirler. Ana plütonik kayalar genellikle olarak Hırfanlı'nın kuzeyinde kuvars diyorit - tonalit, Çelebi - Keskin civarında granodiyorit ve Kırıkkale civarında granit kompozisyonundadır* Buna bağlı olarak da cevherleşme KD yönünde Fe, baz metal ve Mo olarak değişim göstermektedir. Fe cevherleşmeleri içeren Karacaali'deki bazaltlarla, Halildede'deki bazalt ve gabrolar da bu grubun ilk ve en bazik üyeleri olarak kabul edilmektedir. Granodiyorit-granit grubu kayalar KD-GB faylarla bloklara ayrılmış ve bu blokların kuzey yönünde doğuya doğru tedricen ötelenmişler gözlenmiştir. Grubun asidik daykaları genellikle güneyde KB-GD yönünde oldukları halde bu dayklar Kırıkkale'nin doğusunda KD-GB yönünde oluşmuşlardır. Bu yönler aynı zamanda hidrotermal maden yataklarının da esas yönlerini temsil ederler,

Magmatik kayalardaki potasyum miktarının artması ve cevherleşmelerdeki değişimler göz önüne alındığında granodiyorit-granit grubu kayalar ve bunlara bağlı cevherleşmeler iki alt gruba ayrılabilir. Güneybatıdaki birinci alt grup içinde Kesikköprü ve Halildede'de tonalitler Fe, Çelebi'de granodiyorit-granitler W, Karaahmetirde mikrogranitler Mo ve kuzeydoğudaki ikinci alt grup içinde Masatta tonalitler Fe, Keskin'de granodiyorit-granit-mikrogranitler Pb-Zn, Karacaali'de bazaltlar Fe, granit-mikrogranitler Cu-Pb-Zn ve Dağardı ve Balışih'taki granit-mikrogranitler Mo cevherleşmeleri oluşturmuşlardır.

Siyenitler Hamitköy-Âkpinar arasında ve Akçakent yakınlarında mostra verirler. Ayrıca Buzlukdağı ve Çayağzı yakınlarında da ayrı siyenit stokları vardır. Daha genç faz olan kuvars siyenitler florit minerali içerirler ve bölgedeki florit damarlarının oluşumundan da sorumludurlar. Kuvars-florit damarlarının granodiyorit-granit grubu kayalarının içinde bulunması (Küçükbüyük) iki grubun bağlı yaş ilişkilerinin ortaya konulmasında önemli bir rol oynamıştır, İri tane boyulu siyenitler Büyükboyalık yakınlarında K-feldispat açısından değerlendirilebilmektedirler.

Çok alkali kayalar Eşrefi köyünün etrafında dış çapı 4 km'yi bulan karmaşık ring dayk ve stokları şeklinde değişik dokusal özellikler gösteren nefelinli siyenitlerle ve daha çok D-B yönlü ışınal dayklar şeklinde nefelinli siyenit, nefelinli ve lösitli fonolit porfirler ve lösitlerle temsil edilmişlerdir. Akçakentte ayrıca basanit dayklarına da rastlanılmıştır, Alkali kayalar içinde çok sayıda yapı taşı amacına yönelik taş ocağı açılmıştır. Bütün alkali magmatik kayalarda uranyum konsantrasyonu anomali düzeyindedir,

Yozgat kuzeyinde geniş alanlarda mostra veren Tersiyer yaşlı volkanik kayalar Karadere, Sulakyurt ve Yerköy yakınlarında genellikle andezitik ve dasitik lav, tüf ve domları şeklinde görülür. Bu tür kayalar yüksek epitermal cevher potansiyeline sahiptirler. Bölgedeki en genç magmatizma Sulakyurt doğusunda yer alan plato bazaltlarıdır.

IGNEOUS ROCKS AND ORİ DEPOSITS AT THE NORTHERN PART OF KIRŞEHİR

The igneous rocks and related ore deposits at the northern part of Kırşehir were studied in 2001 under the project of MTA, the exploration of mineral deposits related to the acidic magmatic rocks. In addition to the magmatic associations defined at the southern part of Yozgat, five different magmatic associations have been found. From the oldest to the youngest all the associations in the region are as the followings: 1) Meta-basic rocks, 2) Leucogranites, 3) Cefalik dađ granites, 4) Gabbro-basalts, 5) Gabbro-monzonites, 6) Granodiorite-granites, 7) Syenites, 8) Per-alkaline rocks and 9) Tertiary volcanic rocks.

The meta-basic rocks metamorphosed under the green schists faciès are composed of gabbro, basalt, volcano-sedimentary rocks and some acidic volcanic rocks. They are widely outcropped between Balışh and Sulakyurt and include small Fe ores, The leucogranites intrude into the meta-basic rocks along NE-SW extended zones. Generally aplite and pegmatite stock and dykes are less in amount comparing to those in Yozgat, Besides this occurrence, the leucogranites are also seen at the west of Keskin and northeast of Behrekdađh

The Cefalik dađ granites'at the south of Kaman show migmatite character at the south. The mineral distribution within the rock, particularly near the edges of the pluton display very heterogeneous nature in places. The pluton near the edges also shows similar structural patterns with the high degree metamorphic rocks around. The gold occurrences near Savcılıbeyit are thought to be formed by the segregation quartz veins originated during the metamorphism,

The rocks belonged to the gabbro-basalt association occur in the graben basins which pass through at the NW of Kaman with NE direction and at Akçakent with ENE direction. The NW graben near Keskin was not developed very well. Gabbro and monzonite dykes are placed at the peripheries and volcano-sedimentary rocks including basalt and radiolarite at the interior parts of the grabens. Several Mn mine and high Zn anomalies around Kaşımađa are characteristic for this environment. Cutting of the Cefalik dađ granites by the graben and including the leucogranite enclaves by the gabbros of this or next association are evaluated as important features for the relative age relations between the associations.

The gabbro-monzonite association is seen at Baranadađ, Akçakent, and at the south of Dinek dađ. The association shows a normal magmatic zoning, the rocks change inwards from gabbro, diorite, monzodiorite, porphyritic monzonite, to porphyritic quartz monzonite which is similar to the Kerkenez dađ zoning at the south of Sorgun. Dacite dykes and silification together with Fe are sometimes presented in gabbros as those at the south of Bayındır and Akpınar and at Akçakent-Madentepe. The porphyritic monzonites altered by meteoric and/or

55, Türkiye Jeoloji Kurultayı
55th Geological Congress of Turkey

hydrothermal waters are sometimes mined for K-feldspars with a simple enrichment plant such as that in Kilev.

The granodiorite-granite association shows much more different composition at small intervals comparing to that at the south of Yozgat. The main plutonic rocks in general are in composition of quartz diorite-tonolite at the north of Hırfanlı, granodiorite around Çelebi and Keskin and granite around Kırıkkale. Related to this, the ore mineralisation also changes as Fe, base metals and Mo in NE direction. The basalts including Fe at Karacaali and basalts and gabbros near Kesikköprü and Halildede are thought to be the first and most basic members (precursors) of the association. The granodiorite-granites are cut by NE faults and the blocks are pushed towards E in the direction of NE step by step. The acidic members of association generally inclined NW at the south, and NE at the east of Kırıkkale. These also represent the main directions of hydrothermal mineralizations,

When considering the increase in K and changes in ore mineralisation, granodiorite-granite association can be separated into two subassociations. In the first subassociation at the southwest, Fe formed with tonolites in Kesikköprü and Halildede, W with granodiorite-granites in Çelebi and Mo with microgranites in Karahmetli and in the second subassociation at the northeast Fe is together with tonalites in Masat, Pb-Zn with granodiorite-granite-microgranites in Keskin, Fe with basalts and Cu-Pb-Zn with granite-microgranites in Karacaali and Mo with granite-microgranites in Dağardı and Baişlı,

The syenites outcrop between Hamitköy-Akpınar and near Akçakent. There are also some separate syenite stocks at Buzlukdağ and Çayağzı. The younger members of the association are in quartz syenite composition and include disseminated fluorite. This late phase is also responsible for the formation of the quartz-fluorite veins in the area. The presence of the quartz-fluorite veins in granodiorite-granite association (Küçükbüyük) is interpreted as an important evidence for the relative ages between two associations. Coarse grained syenites have been mined for K-feldspar in Büyük Boyalık.

The peralkaline rocks are mainly represented by the nepheline syenites showing various textures and occurring as complex ring dykes and stocks with an outer diameter of 4 km around Eşrefii. The radial dykes extended usually E-W are in composition of nepheline syenite, phonolite porphyries with nepheline and leucite. In addition to these, some basanite dykes occur at Akçakent. Several quarries operate in the peralkaline association for building stone. All the alkaline magmatic rocks show high uranium concentration.

The Tertiary volcanic rocks outcropping in wide areas at the north of Yozgat generally occur around Karadere, Sulakyurt and Yerköy as andesitic and dacitic lavas, tuffs and domes. This kind of rocks own big potential for epithermal mineralisation. The youngest magmatic rocks in the region are the continental plateau basalts at the east of Sulakyurt,